# FERRITIN CONTROL SET

Va	lic	i	y
	Ģ		n

REF: VT-041	Ferritin Control Level I	5 x 1 mL
REF: VT-042	Ferritin Control Level II	5 x 1 mL

Changes made in the instructions for use are marked as grey.

Package insert instructions must be carefully followed. Reliability of assay results cannot be guaranteed if there are any deviations from the instructions in this package.

#### INTENDED USE

Ferritin Control Level I and Ferritin Control Level II are for quality control of Ferritin assay.

#### **CONTENTS / MATERIALS PROVIDED**

Validity Ferritin Control Level I **REF: VT-041** Package: 5 x 1 mL Lyophilized

Validity Ferritin Control Level II **REF: VT-042** Package: 5 x 1 mL Lyophilized

For use with: Validity Ferritin Reagent, Validity Ferritin Calibrator.

**VT-041 / VT-042** do not contain the human serum. Sodium Azide (0,09 %) is added as preservative.

## **CONTROL STABILITY**

Temperature-Conditions	Stability	
Unopened at +2/+8°C	Expiry date on the vial.	
Opened and stored at +25°C	7 days	

## **PREPARATION OF CONTROL**

Control is ready for use.

## INDICATIONS OF INSTABILITY OR DETERIORATION

Presence of extreme turbidity or microbial growth may indicate deterioration.

# PRECAUTIONS

Human source material. Treat as potentially infectious material. Each plasma donor used in the preparation of this product has been tested by an FDA-approved method and found negative for the presence of HIV 1/2 HBsAg, HCV, HIV-Ag antibodies. However, none of the known testing methods can offer complete assurance that the hepatitis B virus, Human Immunodeficiency Virus (HIV) or infectious agents are not present. All human-based products should be handled in accordance with Good Laboratory Practice (GLP) principles using appropriate precautions. Safety data sheets are available at www.validity.com.tr or you may contact your local representative.

## WARNINGS

IVD: For in Vitro Diagnostic use only.

Do not use expired reagents.

Reagents with two different lot numbers should not be interchanged.

For professional use.

Follow Good Laboratory Practice (GLP) guidelines. Contains sodium azide.

CAUTION: Human source samples are processed with this product. All human source samples must be treated as potentially infectious materials and must be handled in accordance with OSHA standards.

# Danger

H317	:May cause allergic skin reaction.
------	------------------------------------

#### Precaution

P280	:Use protective gloves / clothes /
	glasses / mask.
P264	:Wash your hands properly after using.
P272	:Contaminated work clothes should not
	be allowed to be used outside of the
	workplace.
Intervention	
P302+P352	:Wash with plenty of water and soap if
	it contacts with skin.
P333+P313	:Seek medical help if it irritates your
	skin or develops rash.
P362+P364	:Remove contaminated clothes and
	wash properly before using.

## Disposal

# FERRITIN CONTROL SET



# En

P501 :Dispose the vials and contents according to the local regulations.

#### REFERENCES

- Burtis CA, Ashwood ER, Bruns DE, editors. Tiets Textbook of Clinical Chemistry and Molecular Diagnostics, 4th ed. St. Louis, MO, Elsevier Saunders; 2006:2263.
- Occupational Safety and Health Standards: bloodborne pathogens. (29 CFR Part 1910.1030). Fed. Register. July 1, 2001; 17:260 – 273.
- Directive 2000/54/EC. Official Journal of the European Communities No. L262 from October 17, 2000.
- 4. Tietz, N. W. (1983) Clinical guide to Laboratory Tests, p384 W.B. Saunders Co., Philadelphia.

Archem Sağlık Sanayi ve Tic. A.Ş. (With official contract based manufacturing agreement with Validity Sağlık Hiz. San. A.Ş. Company) Mahmutbey Mah. Halkalı Cad. No:124 Kat:4 Bağcılar/İstanbul/Turkey Tel: + 90 212 444 08 92 Fax: +90 212 629 98 89 info@archem.com.tr www.archem.com.tr

info@validity.com.tr www.validity.com.tr

CE

