

ZINC CONTROL LEVEL I-II



REF: VT-025 Zinc Control Level I Liquid 5 x 3 mL REF: VT-026 Zinc Control Level II Liquid 5 x 3 mL

Changes made in the instructions for use are marked as arev.

Package insert instructions must be carefully followed. Reliability of assay results cannot be guaranteed if there are any deviations from the instructions in this package.

INTENDED USE

Validity Zinc Control Level I-II are for quality control of the Zinc Liquid assay.

CONTENTS / MATERIALS PROVIDED

Validity Zinc Control Level I

REF: VT-025

Package: 5 x 3 mL Liquid

Validity Zinc Control Level II

REF: VT-026

Package: 5 x 3 mL Liquid

For use with:

Validity Zinc Liquid Reagent Validity Zinc Calibrator

VT-025 / VT-026 do not contain the human serum.

Materials Required But Not Provided:

- 1. Class A volumetric pipette for liquid transfer
- Distilled or deionized water meeting the specifications equivalent to USP (United States Pharmacopeial Convention) purified water.

CONTROL STABILITY

Temperature-Conditions	Stability
Unopened at +2/+8°C	Expiry date on the vial.
Opened and stored at +2/+8°C	45 days

PREPARATION OF CONTROL

Control is ready for use.

INDICATIONS OF INSTABILITY OR DETERIORATION

Presence of extreme turbidity or microbial growth may indicate deterioration.

PRECAUTIONS



Human source material. Treat as potentially infectious material. Each plasma donor used in the preparation of this product has been tested by an FDA-

approved method and found negative for the presence of HIV 1/2 HBsAg, HCV, HIV-Ag antibodies. However, none of the known testing methods can offer complete assurance that the hepatitis B virus, Human Immunodeficiency Virus (HIV) or infectious agents are not present. All human-based products should be handled in accordance with Good Laboratory Practice (GLP) principles using appropriate precautions. Safety data sheets are available at www.validity.com.tr or you may contact your local representative.

WARNINGS

IVD: For in Vitro Diagnostic use only.

Do not use expired reagents.

Reagents with two different lot numbers should not be interchanged.

For professional use.

Follow Good Laboratory Practice (GLP) guidelines.

CAUTION: Human source samples are processed with this product. All human source samples must be treated as potentially infectious materials and must be handled in accordance with OSHA standards.

Danger

H317 : May cause allergic skin reaction.

Precaution

P280 :Use protective gloves / clothes /

glasses / mask.

P264 :Wash your hands properly after using.
P272 :Contaminated work clothes should not

be allowed to be used outside of the

workplace.

Intervention

P302+P352 :Wash with plenty of water and soap if

it contacts with skin.

P333+P313 :Seek medical help if it irritates your

skin or develops rash.

P362+P364 Remove contaminated clothes and

wash properly before using.

Disposal

P501 :Dispose the vials and contents

according to the local regulations.



ZINC CONTROL LEVEL I-II

En

REFERENCES

- Burtis CA, Ashwood ER, Bruns DE, editors. Tiets Textbook of Clinical Chemistry and Molecular Diagnostics, 4th ed. St. Louis , MO, Elsevier Saunders; 2006:2263.
- 2. Occupational Safety and Health Standards: bloodborne pathogens. (29 CFR Part 1910.1030). Fed. Register. July 1, 2001; 17:260–273.
- 3. Directive 2000/54/EC. Official Journal of the European Communities No. L262 from October 17, 2000.
- 4. Tietz, N. W. (1983) Clinical guide to Laboratory Tests, p384 W.B. Saunders Co., Philadelphia.




Archem Sağlık Sanayi ve Tic. A.Ş. (With official contract based manufacturing agreement with Validity Sağlık Hiz. Sanayi A.Ş. Company)

Mahmutbey Mah. Halkalı Cad. No:124 Kat:4 Bağcılar/İstanbul/Turkey

Tel: + 90 212 444 08 92 **Fax:** +90 212 629 98 89

info@archem.com.tr www.archem.com.tr info@validity.com.tr www.validity.com.tr





Rev: V1.0 Date: 12.2023 Zinc Control Level I-II Page 2 / 2